

Perturbative Quantum Gravity Coupled to Particles in $(1+1)$ -Dimensions

R.B. Mann⁽¹⁾, M. B. Young^(1,2)

¹ *Department of Physics & Astronomy, University of Waterloo,
Waterloo, Ontario, N2L 3G1, Canada*

² *Department of Mathematics and Statistics, Queen's University at Kingston,
Kingston, Ontario, K7L 3N6, Canada*

mann@avatar.uwaterloo.ca, 3my@qmlink.queensu.ca

Abstract

We consider the problem of $(1+1)$ -dimensional quantum gravity coupled to particles. Working with the canonically reduced Hamiltonian, we obtain its post-Newtonian expansion for two identical particles. We quantize the $(1+1)$ -dimensional Newtonian system first, after which explicit energy corrections to second order in c^{-1} are obtained. We compute the perturbed wavefunctions and show that the particles are bound less tightly together than in the Newtonian case.

1 Introduction

It is safe to say that we are still a considerable distance from a satisfactory theory of quantum gravity. Pursuit of this goal is taking place on many fronts, one of which involves working in a two dimensional setting. This furnishes a dynamically simplified framework for developing methods that might be generalizable to higher dimensional quantum gravity, as well as yielding potential insight into quantization of a restricted number of degrees of freedom of such theories.

Although the Einstein vacuum field equations in two spacetime dimensions are trivial insofar as they are satisfied by any metric, it is nevertheless possible to construct dynamically

interesting theories in this setting. This can be done by coupling a scalar field (called the dilaton) into the theory and including suitable matter sources, which typically include the dilaton itself. This has the unfortunate drawback that the theory under consideration typically does not resemble general relativity in its lower-dimensional setting, although some two-dimensional theories are equivalent to a reduction of higher-dimensional Einstein gravity to spherical symmetry [1]. However in their own two-dimensional context, such theories in general do not have a sensible Newtonian limit and/or post-Newtonian expansion [2].

However it has been shown that one can take the $d \rightarrow 2$ limit of general relativity by suitably rescaling Newton's constant G [3]. The resultant theory is one in which the Ricci scalar R is set equal to the trace of the stress energy of prescribed matter fields and sources. In this sense the theory models (3+1)-dimensional general relativity in a manner not captured by other dilaton theories of gravity: the evolution of space-time curvature is governed by the matter distribution, which in turn is governed by the dynamics of space-time [4]. Referred to as $R = T$ theory, when the stress energy is that of a cosmological constant, it reduces to Jackiw-Teitelboim theory [5].

An important feature of the $R = T$ theory is that it has a consistent nonrelativistic ($c \rightarrow \infty$) limit [4] that yields (1+1)-dimensional Newtonian gravity. Its quantization is therefore of considerable potential interest insofar as its (quantum) Newtonian limit can indirectly be empirically tested. This is because the effective theory of gravity near the earth's surface is (1+1)-dimensional Newtonian gravity, whose potential increases linearly with separation away from the surface. From early experimental tests indicating that spin-1/2 particles undergo gravity-induced phase shifts [6] to more recent work demonstrating that neutrons experience a quantization of energy levels in this potential [7], 2D Newtonian quantum gravity affords an important empirical window into semiclassical quantum gravity. In attempting to understand (at least qualitatively) how relativistic effects could modify this behaviour the $R = T$ theory can therefore play an important role.

In this paper we take the first steps toward quantization of the $R = T$ theory. We proceed by taking as a matter source a minimally coupled pair of point particles, forming a 2-body relativistic self-gravitating system. Nonrelativistic (1+1)-dimensional self-gravitating systems (OGS) of N particles have been very important in the fields of astrophysics and cosmology for over 30 years [8], since they not only furnish prototypes for studying gravity in higher dimensions, but also provide effective descriptions of physical systems whose dynamics are closely approximated by the one dimensional system. The OGS phase space is known to exhibit solutions corresponding to very long-lived core-halo configurations, reminiscent

of structures observed in globular clusters [9]. The OGS also approximates collisions of flat parallel domain walls moving in directions orthogonal to their surfaces as well as the dynamics of stars in a direction perpendicular to the plane of a highly flattened galaxy. Relativistic self-gravitating systems have only been more recently studied, primarily in the 2-body [10, 11] and 3-body [12, 13, 14, 15] cases.

In the 2-body case a broad class of exact solutions have been obtained, with extensions that include electromagnetism and cosmological constant [16]. The Hamiltonian is explicitly known as a function of the canonical position and momentum variables. As such its quantization would yield a version of quantum gravity coupled to matter – perhaps the simplest non-trivial quantum gravity system – that affords direct comparison to a classical counterpart. However the nonlinearities of the Hamiltonian yield significant challenges, eg. operator-ordering problems. Here we proceed by quantizing the post-Newtonian expansion of $R = T$ theory. The post-Newtonian expansion for the N -body system has been previously carried out [17]. We consider this expansion for $N = 2$, regarding the post-Newtonian terms as perturbative corrections to (1+1) Newtonian gravity. We find that relativistic corrections tend to lower the ground state energy but raise the energy of all excited states. Furthermore, such corrections tend to spread the wavefunction, indicating a less tightly bound system than in the nonrelativistic case.

The outline of our paper is as follows. We begin with a short review of the relativistic N -body system, outlining briefly how the exact Hamiltonian for the relativistic 2-body system is obtained. We then discuss the post-Newtonian expansion of this Hamiltonian. In section 3 we review the quantization of the non-relativistic system and then in section 4 obtain the post-Newtonian corrections to this system. In section 5 we consider the limits of validity of this approximation, and in section 6 finish off with some concluding remarks. An appendix contains some technical details associated with the perturbative calculations.

2 $R = T$ Gravity and its Post-Newtonian Approximation

We first provide a quick review of the canonical reduction of $(1 + 1)$ -dimensional dilaton gravity. For the gravitational field coupled with N point particles, the action is [17]

$$I = \int dx^2 \left[\frac{1}{2\kappa} \sqrt{-g} \left(\Psi R + \frac{1}{2} g^{\mu\nu} \nabla_\mu \Psi \nabla_\nu \Psi \right) - \sum_{i=1}^N m_i \int d\tau_i \left(-g_{\mu\nu}(x) \frac{dz_i^\mu}{d\tau_i} \frac{dz_i^\nu}{d\tau_i} \right)^{\frac{1}{2}} \delta^2(x - z_i(\tau_i)) \right]. \quad (1)$$

Here $g_{\mu\nu}$ is the metric with $g := \det g_{\mu\nu}$, Ψ is the dilaton field, R is the Ricci scalar, τ_i is the proper time of the i^{th} particle, and $\kappa := \frac{8\pi G}{c^4}$ is the coupling constant, where G is the gravitational constant in two spacetime dimensions and c is the speed of light. ∇_μ is the covariant derivative naturally associated with the metric $g_{\mu\nu}$, $\nabla_\lambda g_{\mu\nu} = 0$.

From the action (1), the field equations for the $R = T$ theory are derived. Upon canonical reduction of the action and field equations [17], the degrees of freedom of the system are z_i and p_i , the position and canonical momentum of the i^{th} particle, respectively. The Hamiltonian density $\mathcal{H}$ is given by

$$\mathcal{H} = -\frac{1}{\kappa} \Delta \Psi$$

where $\Delta = \frac{\partial^2}{\partial x^2}$ is the spatial Laplace operator, and where Ψ is to be viewed as a function of z_i and p_i , determined from the constraint conditions [17]

$$\Delta \Psi - \frac{1}{4} \left(\frac{\partial \Psi}{\partial x} \right)^2 + \kappa^2 \pi^2 + \kappa \sum_i \sqrt{p_i^2 + m_i^2} \delta(x - z_i) = 0 \quad (2)$$

and

$$2 \frac{\partial \pi}{\partial x} + \sum_i p_i \delta(x - z_i) = 0 \quad (3)$$

with π the conjugate momentum to g_{11} . From this, the reduced Hamiltonian of the system is then

$$H = \int dx \mathcal{H}.$$

Once the constraint equations (2) and (3) are solved, H can then be written in terms of z_i and p_i . Since the centre of inertia momentum is conserved and its conjugate variable does not appear in the Hamiltonian, the system has only two degrees of freedom: the separation $r := z_1 - z_2$ and the canonical momentum p of the particles. Solving equations (2) and (3)




Figure 1: The figure shows a plot of both real branches of the Lambert W function, W_0 (solid) and W_{-1} (dot).

for two identical particles of mass m in the centre of inertia frame gives the Hamiltonian implicitly in terms of these variables

$$\left(H - \sqrt{p^2 + m^2} - \epsilon \tilde{p}\right)^2 = \left(\sqrt{p^2 + m^2} - \epsilon \tilde{p}\right)^2 \exp\left(\frac{\kappa}{4}(H - 2\epsilon \tilde{p})|r|\right) \quad (4)$$

where $\tilde{p} := \text{sgn}(r)p$ and ϵ is a constant satisfying $\epsilon^2 = 1$ so as to make time reversal invariance evident.

Eq. (4) can be solved explicitly for H giving [10]

$$H = \sqrt{p^2 c^2 + m^2 c^4} + \text{sgn}(r)pc - \frac{c^4}{\pi G|r|} W(-se^s) \quad (5)$$

where

$$s = \frac{\pi G}{c^4} \left(|r|\sqrt{p^2 c^2 + m^2 c^4} - rpc\right). \quad (6)$$

Note that we have used the explicit form of κ and now show the explicit dependence on c . Here, $W(x)$ is the Lambert W function [18], defined as the solution $W = y(x)$ to the equation $ye^y = x$, which has real solutions for $x \geq -\frac{1}{e}$. W is in general a complex, multivalued function. For $x \in [-\frac{1}{e}, 0)$, W has two real branches. The first, denoted by W_{-1} , satisfies $W(x) \leq -1$. The latter, which is defined for $x \in [-\frac{1}{e}, 0)$, satisfies $-1 \leq W(x)$, is denoted by W_0 and is called the principal branch. See Figure 1 for a plot of both branches. We will always be dealing with the principal branch W_0 , and will simply write for this W . As suggested in Figure 1, we continue W_0 to $x \in [-\frac{1}{e}, \infty)$.

The Hamiltonian (5) is exact to all orders in the gravitational coupling constant $\kappa = \frac{8\pi G}{c^4}$, meaning it accurately describes the system in both the strong and weak field regimes. Expanding it in powers of c^{-1} yields the post-Newtonian approximation¹

$$H = 2mc^2 + \frac{p^2}{m} + 2\pi Gm^2|r| - \frac{4\pi Gm}{c}pr - \frac{p^4}{4m^3c^2} + \frac{4\pi^2 G^2 m^3}{c^2}r^2 + \frac{4\pi G}{c^2}|r|p^2 + \mathcal{O}\left(\frac{1}{c^3}\right). \quad (7)$$

This gives an approximation of Eq. (5) in the case of slow motion in a weak gravitational field. The Hamiltonian (7) contains a term proportional to c^{-1} , which can be removed by making the coordinate transformation [17]

$$r \mapsto \tilde{r}$$

$$pc \mapsto \tilde{p}c + 2\pi Gm^2\tilde{r}.$$

The Hamiltonian (7) then becomes

$$\tilde{H} = \frac{\tilde{p}^2}{m} + 2\pi Gm^2|\tilde{r}| - \frac{\tilde{p}^4}{4m^3c^2} + \frac{4\pi G}{c^2}|\tilde{r}|\tilde{p}^2 + \mathcal{O}\left(\frac{1}{c^3}\right). \quad (8)$$

The above transformation is easily seen to be canonical by the anti-symmetry and linearity of the Poisson brackets, and hence $\tilde{H}(\tilde{r}, \tilde{p})$ plays the rôle of $H(r, p)$ in $(\tilde{r}, \tilde{p})$ space.

In $(3+1)$ -dimensional gravity, terms of odd power in c^{-1} can be associated with gravitational radiation [19]. Because there exists no graviton degree of freedom in the $(1+1)$ -dimensional case, gravitational radiation does not exist, and inclusion of such terms may be misleading. The transformed Hamiltonian $\tilde{H}$ contains relativistic terms proportional only to c^{-2} , more naturally analogous to the $(3+1)$ -dimensional case. From now on the tildes will be dropped for clarity with the understanding that only $(\tilde{r}, \tilde{p})$ coordinates will be used.

If we define $H^0 := \lim_{c \rightarrow \infty} H$, we see that H^0 is the appropriate Hamiltonian to describe two identical particles in $(1+1)$ -dimensional Newtonian gravity. We then have $H = H^0 + H'$ where

$$H^0 = \frac{p^2}{m} + 2\pi Gm^2|r| \quad (9)$$

and

$$H' = -\frac{p^4}{4m^3c^2} + \frac{4\pi G}{c^2}|r|p^2. \quad (10)$$

In the slow moving limit, for small r and p we see $H \approx H^0$, so that the relativistic effects are small compared to the Newtonian effects. This leads to the natural identification of H' as a relativistic perturbation to the Newtonian system.

¹As is common in expansions of relativistic expressions, there is a term proportional to mc^2 that represents the rest energy of the system. This term can be discarded without changing the physics, and hence we will do so from now on.

Note that the Hamiltonian (5) is an exact result that fully encapsulates all of the degrees of freedom of the $R = T$ theory coupled to two point particles. As such, quantization of this Hamiltonian is equivalent to a full quantization of the $R = T$ theory coupled to two point particles in a linear (as opposed to circular) topology. We proceed in the next section to quantize its non-relativistic counterpart (9).

3 Quantization of $(1+1)$ -Dimensional Newtonian Gravity

We would first like to gain an understanding of the two particles in the unperturbed system, described by the Hamiltonian (9). Under the canonical operator identifications in configuration space,

$$r \mapsto r, \quad p \mapsto -i\hbar\partial_r$$

this becomes the Hamiltonian operator

$$\hat{H}^0 = -\frac{\hbar^2}{m}\partial_r^2 + 2\pi Gm^2|r|. \quad (11)$$

The Schrödinger equation then reads $\hat{H}^0\psi_n^0 = E_n^0\psi_n^0$. Here, the subscript labels the wavefunctions and eigenvalues while the superscript identifies them as solutions to the unperturbed (Newtonian) Hamiltonian. Because the potential $V(r) = 2\pi Gm^2|r|$ is continuous on $\mathbb{R}$, non-negative and increases like $|r|$, we know that the system has isolated eigenvalues E_n^0 such that $E_n^0 \rightarrow \infty$ as $n \rightarrow \infty$ [20]. Defining

$$\sigma := \left(\frac{2\pi G}{\hbar^2}\right)^{\frac{1}{3}} m, \quad (12)$$

the normalized wavefunctions are found to be²

$$\psi_n^0(r) = \begin{cases} \sqrt{\frac{\sigma}{2a'_{\frac{n+2}{2}}}} \frac{1}{|Ai(a'_{\frac{n+2}{2}})|} Ai(\sigma|r| + a'_{\frac{n+2}{2}}), & \text{if } n \in \{0\} \cup 2\mathbb{N}, \\ \sqrt{\frac{\sigma}{2}} \frac{1}{|Ai'(a_{\frac{n+1}{2}})|} \text{sgn}(r) Ai(\sigma|r| + a_{\frac{n+1}{2}}), & \text{if } n \in 2\mathbb{N} - 1 \end{cases} \quad (13)$$

with energies

$$E_n^0 = \begin{cases} -(2\pi\hbar G)^{\frac{2}{3}} ma'_{\frac{n+2}{2}}, & \text{if } n \in \{0\} \cup 2\mathbb{N}, \\ -(2\pi\hbar G)^{\frac{2}{3}} ma_{\frac{n+1}{2}}, & \text{if } n \in 2\mathbb{N} - 1 \end{cases} \quad (14)$$

²For the purposes here, an arbitrary phase factor in the normalization constant is of no importance. See Section 4 for details. For normalization procedures involving Airy functions see [21].

and $Ai(x)$ is the Airy function.

In the above, a_n is the n^{th} root of $Ai(x)$, labeled such that $a_{n+1} < a_n$, $n \in \mathbb{N}$, while a'_n is the n^{th} root of $Ai'(x)$, for which an analogous ordering holds (prime denoting the derivative with respect to x). It should be noted that for $n \in \mathbb{N}$, $a_n < a'_n < 0$, so that the energy eigenvalues are strictly positive and monotonically increasing to infinity. The states are non-degenerate, and by the orthonormality relationships $\langle \psi_m^0 | \psi_n^0 \rangle = \delta_{mn}$ [21] and the completeness of $\{\psi_n^0\}_n$ in $L^2(\mathbb{R}, \mathbb{C})$ [22], the eigenfunctions (13) form an orthonormal basis of $L^2(\mathbb{R}, \mathbb{C})$.

4 Quantization of the Post-Newtonian Approximation of $(1 + 1)$ -Dimensional Dilaton Gravity

The Hamiltonian (8) becomes, with the canonical operator identification in configuration space the Hamiltonian operator $\hat{H} = \hat{H}^0 + \hat{H}'$. The spectrum $\{E_n^0\}_n$ and eigenfunctions $\{\psi_n^0\}_n$ of the unperturbed Hamiltonian $\hat{H}^0$ were found explicitly above, and it was noted that the eigenfunctions are non-degenerate and form an orthonormal basis of $L^2(\mathbb{R}, \mathbb{C})$. Hence, the Hamiltonian $\hat{H}$ can be solved approximately using non-degenerate perturbation theory [23]. In the slow moving, weak field limit, we expect the effects of $\hat{H}'$ to be small as compared to those of $\hat{H}^0$. The validity of this assumption will be discussed in Section 5.

For the total Hamiltonian (8), the Schrödinger equation is $\hat{H}\psi_n = E_n\psi_n$. The n^{th} wavefunction ψ_n is written in terms of corrections to the n^{th} Newtonian wavefunction, ψ_n^0 ,

$$\psi_n = \psi_n^0 + \sum_{j \geq 1} \psi_n^j.$$

Here, for $j \in \mathbb{N}$, ψ_n^j is the j^{th} order correction to the unperturbed wavefunction ψ_n^0 . The first order correction is given by [24]

$$\psi_n^1 = \sum_{m \neq n} \frac{\langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle}{E_n^0 - E_m^0} \psi_m^0 \quad (15)$$

where $\langle \cdot | \cdot \rangle : L^2(\mathbb{R}, \mathbb{C}) \times L^2(\mathbb{R}, \mathbb{C}) \rightarrow \mathbb{C}$ is the inner product on the Hilbert space $L^2(\mathbb{R}, \mathbb{C})$ defined by

$$\langle f | g \rangle := \int_{\mathbb{R}} \bar{f} g.$$

In the above, $\bar{f}$ denotes the complex conjugate of f .

The energy eigenvalues are written in an analogous form,

$$E_n = E_n^0 + \sum_{j \geq 1} E_n^j$$

with E_n^j defined as the j^{th} order correction to the unperturbed energy eigenvalue E_n^0 . The first order correction to the energy eigenvalues is given by [24]

$$E_n^1 = \langle \psi_n^0 | \hat{H}' | \psi_n^0 \rangle. \quad (16)$$

We are now in a position to calculate the energy and eigenfunction corrections.

Explicitly, the Hamiltonian operator for the perturbation, $\hat{H}'$, acting on an unperturbed state ψ_n^0 is

$$\hat{H}'\psi_n^0 = -\frac{\hbar^4}{4m^3c^2}\partial_r^4\psi_n^0 - \frac{4\pi\hbar^2G}{c^2}|r|\partial_r^2\psi_n^0 \quad (17)$$

where we have used the so-called natural ordering of operators. Observe that if m is odd and n is even (or vice-versa), we have $\langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle = 0$. Indeed, if n is zero or even, then ψ_n^0 is even, and hence so too is $\partial_r^{2k}\psi_n^0$, $k \in \mathbb{N}$. Then, the inner products $\langle \psi_m^0 | \partial_r^{2k}\psi_n^0 \rangle$ vanish, as they are integrals of an odd function over an even domain. From the definition of ψ_n^1 , we see that this implies that the parity of a state is preserved by the perturbations, as expected since the Hamiltonian (8) has parity symmetry. Hence only when m and n are both odd (or both even) may $\langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle$ be non-vanishing.

We now introduce a useful property of the Airy function that allows the explicit calculation of the desired inner products. It follows from the identity $Ai''(x) = xAi(x)$ that we can write $Ai^{(n)}(x) = f_n(x)Ai(x) + g_n(x)Ai'(x)$, where $f_n(x)$ and $g_n(x)$ satisfy the recursive relationships

$$f_{n+1}(x) = f'_n(x) + xg_n(x), \quad g_{n+1}(x) = f_n(x) + g'_n(x), \quad \forall n \in \mathbb{N} \quad (18)$$

with $f_0(x) = 1$ and $g_0(x) = 0$. For example, using the definitions of f_n and g_n , we get

$$\partial_r^4\psi_n^0 = \sigma^6 r^2 \psi_n^0 + 2\sigma^5 \delta_n |r| \psi_n^0 + \sigma^4 \delta_n^2 \psi_n^0 + 2\sigma^4 \psi_n^{0'} \quad (19)$$

where δ_n is one of a_n or a'_n .

We therefore write for $m, n \in \mathbb{N}$,

$$\begin{aligned} \langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle &= - \left(\frac{\hbar^4}{4m^3c^2} \sigma^6 + \frac{4\pi\hbar^2G}{c^2} \sigma^3 \right) \langle \psi_m^0 | r^2 | \psi_n^0 \rangle \\ &\quad - \left(\frac{\hbar^4}{2m^3c^2} \sigma^5 \delta_n + \frac{4\pi\hbar^2G}{c^2} \sigma^2 \delta_n \right) \langle \psi_m^0 | |r| | \psi_n^0 \rangle \\ &\quad - \frac{\hbar^4}{4m^3c^2} \sigma^4 \delta_n^2 \langle \psi_m^0 | \psi_n^0 \rangle - \frac{\hbar^4}{2m^3c^2} \sigma^4 \langle \psi_m^0 | \psi_n^{0'} \rangle. \end{aligned} \quad (20)$$

We evaluate the above expression with $m = n$, which gives E_n^1 . All inner products can be evaluated explicitly.³ After putting in the explicit form of σ , we obtain

$$E_n^1 = \begin{cases} \frac{m}{2} \frac{(2\pi\hbar G)^{\frac{4}{3}}}{c^2} \left(\frac{13}{30} \left(a'_{\frac{n+2}{2}} \right)^2 + \frac{9}{10} \frac{1}{a'_{\frac{n+2}{2}}} \right) & \text{if } n \in \{0\} \cup 2\mathbb{N}, \\ \frac{m}{2} \frac{(2\pi\hbar G)^{\frac{4}{3}}}{c^2} \left(\frac{13}{30} a_{\frac{n+1}{2}}^2 \right) & \text{if } n \in 2\mathbb{N} - 1. \end{cases} \quad (21)$$

Inserting appropriate values of a_n and a'_n shows that $E_0^0 < 0$ while $E_n^0 > 0$ for $n \geq 1$. By the ordering and unboundedness of a_n and a'_n , E_n^1 is monotonically increasing in n without bound, and so the approximation will be valid for only finitely many states, a point we shall return to in Section 5. The magnitude of the energy corrections is expected to be quite small, as is common in quantum gravity. Indeed, for the ground state, we have

$$\left| \frac{E_0^1}{E_0^0} \right| \approx 0.21 \frac{(2\pi\hbar G)^{\frac{2}{3}}}{c^2}, \quad (22)$$

where the lower dimensionality implies that G has the same dimensions as $c^3/\hbar$.

Proceeding, we can now find expressions for the perturbed wavefunctions. To do so, we evaluate (20) with m and n both zero or even to get

$$\langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle = \frac{m}{2} \frac{(2\pi\hbar G)^{\frac{4}{3}}}{c^2} \left[54 + 5a'_{\frac{n+2}{2}} \left(a'_{\frac{m+2}{2}} - a'_{\frac{n+2}{2}} \right)^2 \right] \frac{(-1)^{m+n+1} \left(a'_{\frac{m+2}{2}} + a'_{\frac{n+2}{2}} \right)}{\left(a'_{\frac{m+2}{2}} - a'_{\frac{n+2}{2}} \right)^4 \sqrt{a'_{\frac{m+2}{2}} a'_{\frac{n+2}{2}}}} \quad (23)$$

and for m and n both odd,

$$\langle \psi_m^0 | \hat{H}' | \psi_n^0 \rangle = \frac{m}{2} \frac{(2\pi\hbar G)^{\frac{4}{3}}}{c^2} \left[54 + 5a_{\frac{n+1}{2}} \left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}} \right)^2 \right] \frac{(-1)^{m+n} 2}{\left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}} \right)^4}. \quad (24)$$

Using Eqs. (15), (23) and (24), we have

$$\psi_n^1 = \sum_{m \geq 0} \alpha_{mn} \psi_m^0 \quad (25)$$

where

$$\alpha_{mn} = \begin{cases} \frac{(2\pi\hbar G)^{\frac{2}{3}}}{2c^2} (-1)^{m+n+1} \frac{\left(54 + 5a'_{\frac{n+2}{2}} \left(a'_{\frac{m+2}{2}} - a'_{\frac{n+2}{2}} \right)^2 \right) \left(a'_{\frac{m+2}{2}} + a'_{\frac{n+2}{2}} \right)}{\left(a'_{\frac{m+2}{2}} - a'_{\frac{n+2}{2}} \right)^5 \sqrt{a'_{\frac{m+2}{2}} a'_{\frac{n+2}{2}}}}, & \text{if } m \neq n, m, n \in \{0\} \cup 2\mathbb{N} \\ \frac{(2\pi\hbar G)^{\frac{2}{3}}}{c^2} (-1)^{m+n} \frac{\left(54 + 5a_{\frac{n+1}{2}} \left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}} \right)^2 \right)}{\left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}} \right)^5}, & \text{if } m \neq n, m, n \in 2\mathbb{N} - 1 \\ 0, & \text{otherwise.} \end{cases} \quad (26)$$

³See Appendix.




Figure 2: The figure to the left shows a plot of ψ_0^0 (solid), ψ_0 for $c = 7$ (dash) and ψ_0 for $c = 3$ (dash-dot). On the right, the plot shows $|\psi_0^0|^2$ (solid), $|\psi_0|^2$ for $c = 7$ (dash) and $|\psi_0|^2$ for $c = 3$ (dash-dot). Units are chosen such that $\hbar = 1J \cdot s$, $G = \frac{1}{2\pi} \frac{m}{kg \cdot s^2}$, and $m = 1kg$.

The first order perturbed wavefunctions are thus given by⁴

$$\psi_n = \psi_n^0 + \sum_{m \geq 0} \alpha_{mn} \psi_m^0. \quad (27)$$

It is easy to check that to second order in c^{-1} , ψ_n is normalized. See Figure 2 for a plot of the perturbed wavefunctions and probability densities for increasing values of c .

5 Expectation Values and the Validity of the Corrections

We now wish to calculate the expectation values of powers of $\hat{r}$ and $\hat{p}$ so as to compare the Newtonian and post-Newtonian systems, and to check the limits of validity of our perturbation results. First, some notation. We define, for $\phi \in D(\hat{Q}) \subset L^2(\mathbb{R}, \mathbb{C})$,

$$\langle \hat{Q} \rangle_\phi := \langle \phi | \hat{Q} | \phi \rangle \quad (28)$$

for some observable $\hat{Q}$.

We state results for the perturbed wavefunctions, which are derived using the inner product identities developed in the Appendix. Taking the $c \rightarrow \infty$ limit, these also give the

⁴Had a phase factor been included in ψ_n^0 , say $\tilde{\psi}_n^0 = e^{i\theta_n} \psi_n^0$ for some $\theta_n \in \mathbb{R}$, then we would now have that $\tilde{\psi}_n = e^{i\theta_n} \psi_n$. For simplicity we omit this phase factor.

Newtonian expectation values. We have, keeping only terms to second order in c^{-1} ,

$$\langle \hat{r} \rangle_{\psi_n} = 0, \quad \langle \hat{p} \rangle_{\psi_n} = 0, \quad (29)$$

reflecting the fact that the perturbation (10) is parity preserving. We also find

$$\langle \hat{r}^2 \rangle_{\psi_n} = \begin{cases} \left[\frac{8a_{\frac{n+2}{2}}'^3 - 3}{15a_{\frac{n+2}{2}}'} + 24 \sum_{m \geq 0} \alpha_{mn} \frac{(-1)^{m+n} \left(a_{\frac{m+2}{2}}' + a_{\frac{n+2}{2}}' \right)}{\left(a_{\frac{n+2}{2}}' - a_{\frac{m+2}{2}}' \right)^4 \sqrt{a_{\frac{m+2}{2}}' a_{\frac{n+2}{2}}'}} \right] \left(\frac{\hbar^2}{2\pi G} \right)^{\frac{2}{3}} \frac{1}{m^2}, & \text{if } n \in \{0\} \cup 2\mathbb{N} \\ \left[\frac{8a_{\frac{n+1}{2}}^2}{15} + 48 \sum_{m \geq 0} \alpha_{mn} \frac{(-1)^{m+n+1}}{\left(a_{\frac{n+1}{2}} - a_{\frac{m+1}{2}} \right)^4} \right] \left(\frac{\hbar^2}{2\pi G} \right)^{\frac{2}{3}} \frac{1}{m^2}, & \text{if } n \in 2\mathbb{N} - 1, \end{cases} \quad (30)$$

and

$$\langle \hat{p}^2 \rangle_{\psi_n} = \begin{cases} - \left[\frac{1}{3} a_{\frac{n+2}{2}}' + 2 \sum_{m \geq 0} \alpha_{mn} \frac{(-1)^{m+n} \left(a_{\frac{m+2}{2}}' + a_{\frac{n+2}{2}}' \right)}{\left(a_{\frac{n+2}{2}}' - a_{\frac{m+2}{2}}' \right)^2 \sqrt{a_{\frac{m+2}{2}}' a_{\frac{n+2}{2}}'}} \right] (2\pi \hbar G)^{\frac{2}{3}} m^2, & \text{if } n \in \{0\} \cup 2\mathbb{N} \\ - \left[\frac{1}{3} a_{\frac{n+1}{2}} + 2 \sum_{m \geq 0} \alpha_{mn} \frac{(-1)^{m+n+1}}{\left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}} \right)^2} \right] (2\pi \hbar G)^{\frac{2}{3}} m^2, & \text{if } n \in 2\mathbb{N} - 1. \end{cases} \quad (31)$$

We see that $\langle \hat{r}^2 \rangle_{\psi_n}$ and $\langle \hat{p}^2 \rangle_{\psi_n}$ depend strongly on the masses of the particles, being proportional to the inverse square and square of the mass, respectively. This dependence on mass resembles an exaggerated version of the simple harmonic oscillator (SHO), with $\langle \hat{r}^2 \rangle^{SHO}$ and $\langle \hat{p}^2 \rangle^{SHO}$ being proportional to the inverse root and root of the mass, respectively. For particles of small (large) mass, the square separation quickly becomes very large (small) and changes slowly (quickly) due to the m^2 dependence of $\langle \hat{p}^2 \rangle_{\psi_n}$. It is also instructive to compare these results to those of $(3+1)$ -dimensional Newtonian gravity. In the $(3+1)$ case, $\langle \hat{r}^2 \rangle$ and $\langle \hat{p}^2 \rangle$ are proportional to m^{-6} and m^6 , respectively. In this sense the expectation values are an even more exaggerated version of the SHO values.

From Eq. (26), we see that $\lim_{c \rightarrow \infty} \alpha_{mn} = 0$. Using this, it is straightforward to check that we have $\lim_{c \rightarrow \infty} \langle \hat{Q} \rangle_{\psi_n} = \langle \hat{Q} \rangle_{\psi_n^0}$ for $\hat{Q} = \hat{r}^n, \hat{p}^n$, $n = 1, 2$, as claimed above. This is illustrated in Figure 3. From the expression for α_{mn} , we see that the expectation values diverge as $c \rightarrow 0$. This, however, is not of concern as the approximation is valid only for large c . Higher order




Figure 3: The figure to the left shows a plots, for $n = 0, 1$, of $\langle \hat{r}^2 \rangle_{\psi_n}$ (dot) as a function of c and $\langle \hat{r}^2 \rangle_{\psi_n^0}$ (solid), illustrating $\lim_{c \rightarrow \infty} \langle \hat{r}^2 \rangle_{\psi_n} = \langle \hat{r}^2 \rangle_{\psi_n^0}$. To the right, plots of $\langle \hat{p}^2 \rangle_{\psi_n}$, $n = 0, 1$, (dot) as a function of c and $\langle \hat{p}^2 \rangle_{\psi_n^0}$ (solid) are shown, illustrating $\lim_{c \rightarrow \infty} \langle \hat{p}^2 \rangle_{\psi_n} = \langle \hat{p}^2 \rangle_{\psi_n^0}$. Units are chosen such that $\hbar = 1J \cdot s$, $G = \frac{1}{2\pi} \frac{m}{kg \cdot s^2}$, and $m = 1kg$.

contributions in c^{-1} will greatly modify Figure 3 for small c , while leaving the behaviour mostly unchanged for large c .

As an example, in units such that $\hbar = 1$, $G = \frac{1}{2\pi}$, and $m = 1$, we have $\langle \hat{r}^2 \rangle_{\psi_0^0} \approx 0.7499$ and $\langle \hat{p}^2 \rangle_{\psi_0^0} \approx 0.3396$. Setting $c = 10$, and including twenty terms (i.e. $m = 0, \dots, 19$) in the sums, we get $\langle \hat{r}^2 \rangle_{\psi_0} \approx 0.7636$ and $\langle \hat{p}^2 \rangle_{\psi_0} \approx 0.3341$.

Denoting the dispersion of an observable $\hat{Q}$ in the state ϕ by $(\Delta \hat{Q})_\phi$, using the above expectation values, we see that

$$(\Delta \hat{x})_{\psi_n^0} (\Delta \hat{p})_{\psi_n^0} = \begin{cases} \frac{1}{3} \sqrt{\frac{3-8 \left(\frac{a'_{n+2}}{5} \right)^3}{5}} \hbar, & \text{if } n \in \{0\} \cup 2\mathbb{N} \\ \frac{2}{3} \sqrt{\frac{2}{5}} |a_{\frac{n+1}{2}}| \hbar, & \text{if } n \in 2\mathbb{N} - 1. \end{cases} \quad (32)$$

It is straightforward to show that the uncertainty principle is not violated upon substituting values for a_n and a'_n . Figure 4 shows a plot of $(\Delta \hat{x})(\Delta \hat{p})$ for both the perturbed and unperturbed ground state wavefunctions, as well as the lower bound $\frac{\hbar}{2}$ set by the uncertainty principle. Of course for small enough c we find that $(\Delta \hat{x})_{\psi_0^0} (\Delta \hat{p})_{\psi_0^0} < \frac{\hbar}{2}$, indicative of the limits of the validity of our approximation.

To obtain an estimate of the number of states for which our approximation holds, we demand that the expected gravitational binding energy be less than the rest energy of the system [19]:

$$2\pi G m^2 |r| < 2mc^2.$$




Figure 4: The figure shows a plot $\frac{\hbar}{2}$ (solid), $(\Delta\hat{x})_{\psi_0^0}(\Delta\hat{p})_{\psi_0^0}$ and $(\Delta\hat{x})_{\psi_0}(\Delta\hat{p})_{\psi_0}$ (both dash) as a function of c , illustrating $\lim_{c \rightarrow \infty} (\Delta\hat{x})_{\psi_0}(\Delta\hat{p})_{\psi_0} = (\Delta\hat{x})_{\psi_0^0}(\Delta\hat{p})_{\psi_0^0}$. Units are such that $\hbar = 1J \cdot s$, $G = \frac{1}{2\pi} \frac{m}{kg \cdot s^2}$, and $m = 1kg$.

Quantum mechanically, this suggests the bound

$$\langle \hat{r}^2 \rangle_{\psi_n} < \frac{c^4}{\pi^2 G^2 m^2}$$

which is a bound on n , since it requires

$$\frac{8a_{\frac{n+1}{2}}^2}{15} + 48 \sum_{m \geq 0} \alpha_{mn} \frac{(-1)^{m+n+1}}{\left(a_{\frac{n+1}{2}} - a_{\frac{m+1}{2}}\right)^4} < \frac{4c^4}{(2\pi\hbar G)^{\frac{4}{3}}}. \quad (33)$$

For $\hbar = 1$, $G = \frac{1}{2\pi}$, $c = 10$, this gives $n \lesssim 645$. So, although the number of states is finite, even for small c , this number is quite large.

6 Discussion

As noted above, the Hamiltonian (5) represents the exact energy functional of all of the degrees of freedom in the relativistic 2-body system. Hence its quantization is tantamount to a full quantization of gravity coupled to matter in (1+1)-dimensions.

We have successfully completed the first steps to this end. We have used perturbation theory to obtain explicit energy and eigenfunction corrections to the relativistically perturbed Newtonian Hamiltonian to second order in c^{-1} . We found that the ground state energy in the

relativistic case is lower than in the Newtonian case, while all other relativistic states have higher energies than their Newtonian counterparts. As suggested in Figure 3, we conjecture that $\langle \hat{r}^2 \rangle_{\psi_n^0} < \langle \hat{r}^2 \rangle_{\psi_n}$ and $\langle \hat{p}^2 \rangle_{\psi_n^0} > \langle \hat{p}^2 \rangle_{\psi_n}$. That is, relativistically, the two particles are not bound as tightly together as in Newtonian theory. The above inequalities have been verified numerically for $n \leq 20$, including forty terms in the sums. On the other hand, on average we expect the separation of the particles to be changing more slowly in the relativistic case than in its Newtonian counterpart.

In estimating the actual magnitude of the quantum-gravitational corrections to the Newtonian system, we run into the problem of not knowing the appropriate values of the fundamental constants c , $\hbar$ and G in the $(1+1)$ -dimensional case. However if we view the $R = T$ theory as being the effective relativistic theory near the surface of the earth we can obtain a numerical value of the energy correction. The $(3+1)$ Newtonian potential close to the surface of the Earth is

$$V^{(4)}(r) = -\frac{G^{(4)} M_{\oplus} m}{|R_{\oplus} + r|}.$$

For small r , this becomes

$$V^{(4)}(r) = -\frac{G^{(4)} M_{\oplus} m}{R_{\oplus}} + \frac{G^{(4)} M_{\oplus} m}{R_{\oplus}^2} r + \mathcal{O}(r^2).$$

On the other hand, the $(1+1)$ Newtonian potential for the two masses is

$$V^{(2)}(r) = 2\pi G M_{\oplus} m |r|.$$

Here the superscripts in parentheses refer to the spacetime dimension in which the quantity is valid (where $G := G^{(2)}$). Ignoring the constant term, comparison of the two cases suggests

$$G = \frac{G^{(4)}}{2\pi R_{\oplus}^2} \tag{34}$$

which indeed provides G with the correct units. Using this, with the usual values of $\hbar$ and c , we find $\left| \frac{E_1^1}{E_0^0} \right| \approx 8 \times 10^{-56}$.

Of course the actual theory correcting Newtonian gravity in $(3+1)$ dimensions is general relativity and not the $R = T$ theory. However the results obtained here illustrate – quantitatively and qualitatively – that sensible quantum-gravitational corrections to Newtonian gravity can be meaningfully computed, though the suppression of the effective 2-dimensional Newton constant G renders such effects unobservable in practice. A full quantization of the $R = T$ theory remains an interesting problem for future study.

7 Appendix

7.1 Inner products of the form $\langle \psi_m^0 || r^i | \psi_n^0 \rangle$

In calculating inner products involving ψ_n^0 , we are led to consider integrals of the form $\int_0^\infty dr |r|^i Ai(r + \delta_1) Ai(r + \delta_2)$, $i = 1, 2$ and $\delta_j \in \mathbb{R}$. Such integrals are found in [21]. We must investigate the cases in which $\delta_1 = \delta_2$ and $\delta_1 \neq \delta_2$ separately.

7.1.1 $\delta_1 = \delta_2$

We will show explicit calculations for the case $q = 1$, which illustrates the general procedure, and state the remaining cases as results. From [21],

$$\begin{aligned} \int dx x Ai^2(x + \delta) &= \frac{1}{3} (x^2 - x\delta - 2\delta^2) Ai^2(x + \delta) + \frac{2\delta - x}{3} Ai'^2(x + \delta) \\ &\quad + \frac{1}{3} Ai'(x + \delta) Ai(x + \delta). \end{aligned}$$

Using this result for ψ_n^0 , with, say, n odd, we get after setting $\rho := \sigma r$,

$$\langle \psi_n^0 || r | \psi_n^0 \rangle = \frac{2a_{\frac{n+1}{2}}^2}{\sigma^2} \int_0^\infty d\rho \rho Ai^2\left(\rho + a_{\frac{n+1}{2}}\right) = -\frac{4a_{\frac{n+1}{2}}^2}{3\sigma^2} Ai'^2\left(a_{\frac{n+1}{2}}\right).$$

Using the expression for a_n , we see get the final result⁵

$$\langle \psi_n^0 || r | \psi_n^0 \rangle = -\frac{2a_{\frac{n+1}{2}}}{3\sigma}.$$

Similarly, for n even, we get

$$\langle \psi_n^0 || r | \psi_n^0 \rangle = -\frac{2a'_{\frac{n+2}{2}}}{3\sigma}.$$

For n odd,

$$\langle \psi_n^0 | r^2 | \psi_n^0 \rangle = \frac{8a_{\frac{n+1}{2}}^2}{15\sigma^2},$$

while for n even,

$$\langle \psi_n^0 | r^2 | \psi_n^0 \rangle = \frac{8a_{\frac{n+2}{2}}'^3 - 3}{15\sigma^2 a'_{\frac{n+2}{2}}}.$$

⁵We used the fact that $Ai(x) \sim \frac{1}{2\sqrt{\pi}} \frac{e^{-\frac{2}{3}x^{\frac{3}{2}}}}{x^{\frac{1}{4}}}$ as $x \rightarrow \infty$ [25], so that $x^q Ai(x) \rightarrow 0$ as $x \rightarrow \infty \forall q \in \mathbb{N}$.

7.1.2 $\delta_1 \neq \delta_2$

Again, an explicit calculation will be shown for the case $q = 1$ case, while the remaining cases will be stated without derivations. From [21],

$$\begin{aligned} \int dx x Ai(x + \delta_1) Ai(x + \delta_2) &= -\frac{\delta_1 + \delta_2 + 2x}{(\delta_1 - \delta_2)^2} Ai(x + \delta_1) Ai(x + \delta_2) \\ &+ \left[\frac{x}{\delta_1 - \delta_2} + \frac{2}{(\delta_1 - \delta_2)^3} \right] \{ Ai(x + \delta_1) Ai'(x + \delta_2) \\ &- Ai'(x + \delta_1) Ai(x + \delta_2) \} \\ &+ \frac{2}{(\delta_1 - \delta_2)^2} Ai'(x + \delta_1) Ai'(x + \delta_2). \end{aligned}$$

Using this result for ψ_n^0 , with, say, m and n both odd, we get after setting $\rho := \sigma r$,

$$\langle \psi_m^0 | r | \psi_n^0 \rangle = -\frac{4\bar{a}_{\frac{m+1}{2}} a_{\frac{n+1}{2}} Ai'\left(a_{\frac{m+1}{2}}\right) Ai'\left(a_{\frac{n+1}{2}}\right)}{\sigma^2 \left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}}\right)}.$$

Using the expression for a_n , we see get the final result

$$\langle \psi_m^0 | r | \psi_n^0 \rangle = \frac{(-1)^{m+n+1} 2}{\sigma \left(a_{\frac{m+1}{2}} - a_{\frac{n+1}{2}}\right)^2}.$$

For m and n both even, we have

$$\langle \psi_m^0 | r | \psi_n^0 \rangle = \frac{(-1)^{m+n} (a'_{\frac{m+2}{2}} + a'_{\frac{n+2}{2}})}{\sigma \left(a'_{\frac{m+2}{2}} - a'_{\frac{n+2}{2}}\right)^2 \sqrt{a'_{\frac{m+2}{2}} a'_{\frac{n+2}{2}}}}.$$

For m and n both odd, we have

$$\langle \psi_m^0 | r^2 | \psi_n^0 \rangle = \frac{(-1)^{m+n+1} 24}{\sigma^2 \left(a_{\frac{n+1}{2}} - a_{\frac{m+1}{2}}\right)^4}.$$

For m and n both even, we have

$$\langle \psi_m^0 | r^2 | \psi_n^0 \rangle = \frac{(-1)^{m+n} 12 \left(a'_{\frac{m+2}{2}} + a'_{\frac{n+2}{2}}\right)}{\sigma^2 \left(a'_{\frac{n+2}{2}} - a'_{\frac{m+2}{2}}\right)^4 \sqrt{a'_{\frac{m+2}{2}} a'_{\frac{n+2}{2}}}}.$$

Acknowledgement

This work was supported by the Natural Sciences & Engineering Research Council of Canada.

References

- [1] D. Grumiller, W. Kummer and D. V. Vassilevich, Phys. Rept. **369**, 327 (2002) (hep-th/0204253).
- [2] S.F.J. Chan and R.B. Mann, Class. Quant. Grav. **12**, 351(1995).
- [3] R.B. Mann and S.F. Ross, Class.Quant.Grav. **10**, 345 (1993).
- [4] R.B. Mann, Found. Phys. Lett. **4** (1991) 425; R.B. Mann, Gen. Rel. Grav. **24** (1992) 433.
- [5] R. Jackiw, Nucl. Phys. B **252**, 343 (1985); C. Teitelboim, Phys. Lett. B **126**, 41, (1983).
- [6] R. Colella, A. W. Overhauser and S. A. Werner, Phys. Rev. Lett. **34**, 1472 (1975).
- [7] V. Nesvizhevsky et. al. Nature **415**, 297 (2001).
- [8] G. Rybicki, Astrophys. Space. Sci **14** (1971) 56; H.L. Wright, B.N. Miller, and W.E. Stein, Astrophys. Space. Sci. **84**, **2** (1982) 421 and references therein.
- [9] See B.N. Miller and P. Youngkins, Phys. Rev. Lett. **81** 4794 (1998); K.R. Yawn and B.N. Miller, Phys. Rev. Lett. **79** 3561 (1997) and references therein.
- [10] R.B. Mann and T. Ohta, Phys. Rev. **D55**, 4723 (1997), gr-qc/9611008; Class. Quant. Grav. **14**, 1259 (1997).
- [11] R.B. Mann, D. Robbins and T. Ohta, Phys.Rev.Lett. **82**, 3738 (1999).
- [12] F. Burnell, R.B. Mann and T. Ohta, “Chaos in a Relativistic 3-body Self-Gravitating System”, Phys. Rev. Lett. **90** (2003) 134101.
- [13] F.J. Burnell, J.J. Malecki, R.B. Mann, and T. Ohta, “Chaos in an Exact Relativistic 3-body Self-Gravitating System”, Phys. Rev. **E69** (2004) 016214 .
- [14] J. J. Malecki and R. B. Mann, Phys. Rev. **E69** (2004) 066208.
- [15] S. Bachmann, M.J. Koop and R.B. Mann , astro-ph/0607070

- [16] R.B. Mann, D. Robbins, T. Ohta and M. Trott, Nucl. Phys. **B590** 367 (200).
- [17] R.B. Mann and T. Ohta, Class. Quantum Grav. **13** 2585 (1996), gr-qc/9605004.
- [18] R.M. Corless *et al.*, Adv. Comp. Math. **5** 329 (1996).
- [19] S. Weinberg. *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*. John Wiley & Sons, Inc., New York, N.Y. 1972.
- [20] S. Gustafson and I. Sigal. *Mathematical Concepts in Quantum Mechanics*. Springer, New York, N.Y. 2003.
- [21] O. Vallée and M. Soares. *Airy Functions and Applications to Physics*. Imperial College Press, London, U.K. 2004.
- [22] E.C. Titchmarsh. *Eigenfunction Expansions Associated with Second-Order Differential Equations*. Oxford University Press, Oxford, U.K. 1962.
- [23] F.M. Fernández. *Introduction to Perturbation Theory in Quantum Mechanics*. CRC Press, Boca Raton, Fl. 2001.
- [24] D.J. Griffiths. *Introduction to Quantum Mechanics*. Prentice Hall, Upper Saddle River, N.J. 2005.
- [25] M. Abramowitz and I. Stegun. *Handbook of Mathematical Functions*. Dover Publications, New York, N.Y. 1965.